

CORRESPONDENCE

Is It Right ASA Scoring to be Used in Identification of Nonoperated Patients?

Although with limited number of patients, we appreciate the authors for their study investigating the rates of rupture and mortality that was developed in inoperable abdominal aortic aneurysm.¹ First we wanted to state a technical mistake in the paper that drew our attention. Despite the total number of patients included in the study is 72, there are 71 patients in Figure 1. Since the number of patients is small, we believe that this missing patient should be included in the relevant group. However, our actual review about the article is that the authors grouped the patients according to the ASA and gave the number of deaths and ruptures of these groups. Several studies reported ASA scoring to be effective both on anesthetic and surgical outcome.^{2,3} However, we believe that this parameter which is used in preoperative risk scoring should not be used to identify nonoperated patients. Already looking to Figure 1, survival rate in the ASA 4 group (48%) is seen to be unexpectedly higher than ASA 2 group (38%). This shows us that evaluation of the risk for rupture and mortality between ASA groups is meaningless. We would want to say that we wonder about the views of the authors on our this critics.

REFERENCES

- 1 Western CE, Carlisle J, McCarthy RJ, Currie IC. Palliation of abdominal aortic aneurysms in the endovascular era. *Eur J Vasc Endovasc Surg* 2012 Nov 14.
- 2 Wisniewski B, Barnes M, Jenkins J, Boyne N, Kruger A, Walker PJ. Predictors of outcome after elective endovascular abdominal aortic aneurysm repair and external validation of a risk prediction model. *J Vasc Surg* 2011 Sep;54(3):644–53.
- 3 Barnes M, Boulton M, Maddern G, Fitridge R. A model to predict outcomes for endovascular aneurysm repair using preoperative variables. *Eur J Vasc Endovasc Surg* 2008 May;(5):571–9.

O. Gokalp*

*Izmir Katip Celebi University, Faculty of Medicine,
Department of Cardiovascular Surgery, Turkey*

I.S. Satoglu

*Izmir Katip Celebi University, Faculty of Medicine,
Department of Orthopaedics and Traumatology, Turkey*

Y. Besir, H. Cakir

*Izmir Katip Celebi University, Atatürk Education and
Research Hospital, Department of Cardiovascular Surgery,
Turkey*

H. Kara

*Giresun Ada Hospital, Department of Cardiovascular
Surgery, Turkey*

A. Gurbuz

*Izmir Katip Celebi University, Faculty of Medicine,
Department of Cardiovascular Surgery, Turkey*

*Corresponding author. O. Gokalp, Altinvadi Cd. No:85,
D:10 35320 Narlidere, Izmir, Turkey. Tel.: +90 5052168813;
fax: +90 2322431530.

Email-addresses: gokalporhan@yahoo.com,
gokalporhan@myinet.com (O. Gokalp)

Available online 6 February 2013

© 2013 European Society for Vascular Surgery. Published by
Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.ejvs.2013.01.014>

Re. 'Is It Right ASA Scoring to be Used in Identification of Nonoperated Patients?'

Sir,

We thank Dr Gokalp and colleagues for their interest in our paper on palliation of aortic aneurysms. Our study contained 72 patients however, we were unable to determine cause of death in one individual as they died abroad and therefore we were unable to include them in the analysis of cause of death.

ASA grade has been used in many studies to stratify surgical risk.^{1,2} ASA grade is only one factor considered in our unit when deciding suitability for aneurysm repair and, unsurprisingly, the majority of patients included in this study were ASA-3 (41) or ASA-4 (22). Of the small number of ASA-2 patients (8), five declined operative intervention, some of whom would have been suitable candidates for aneurysm repair, so possibly changing subsequent outcome (25% of this cohort died of rupture). We therefore feel that as the majority of this group in our study self-selected themselves out of surgery, they cannot be seen as representative in terms of outcome.

Following assessment by a Consultant Vascular Surgeon, patients in our unit are subjected to CPEX testing and Consultant Anaesthetist review. This was not consistent in the early days of our CPEX programme but is standard practice now. All these factors are combined to give a definitive judgement on suitability for intervention at this point in time. This can be revisited in the future if aneurysm expansion shifts risk-benefit ratio in favour of surgery.

REFERENCES

- 1 Lomazzi C, Mariscalco G, Piffaretti G, Bacuzzi A, Tozzi M, Carrafiello G, et al. Endovascular treatment of elective abdominal aortic aneurysms: independent predictors of early and late mortality. *Ann Vasc Surg* 2011;25(3):299–305.
- 2 Zannetti S, DeRango P, Parlani G, Verzini F, Maselli A, Cao P. Endovascular abdominal aortic aneurysm repair in high-risk patients: a single centre experience. *Eur J Vasc Endovasc Surg* 2001;21(4):334–8.

C.E. Western*

*Torbay Hospital, Vascular Surgery, Lawes Bridge, Torquay,
Devon TQ2 2AA, United Kingdom*

*Tel.: +44 7788547537.

Email-address: catherinewestern@yahoo.co.uk

Available online 19 February 2013

© 2013 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.ejvs.2013.01.017>

DOI of original article: <http://dx.doi.org/10.1016/j.ejvs.2013.01.014>

Re. 'An Online Patient Completed Aberdeen Varicose Vein Questionnaire Can Help to Guide Primary Care Referrals'

We read with interest the study by Ward et al. concerning an online completion of the disease specific quality of life Aberdeen Varicose Vein questionnaire (AVVQ).¹ The study is thought provoking however surely the use of a non-validated tool renders such extrapolations moot. Without the original full questionnaire no comparisons can be drawn. Multiple other questionnaires exist which do not require a drawing such as Chronic Lower Limb Venous Insufficiency Questionnaire (CIVIQ)² or Specific Quality of Life and Outcome Response – Vascular (SQOR-V).³ Additionally, construction of an online drawing tool is now well within our technological grasp.

CEAP grade 4–6 showed a high prevalence in this study (42%) with a consequently high average VCSS (mean 18.6). However, this is not reflected in the AVVQ scores (mean 21.8). Additionally the correlation for CEAP and AVVQ scores is not documented – is this clinically as well as statistically significant? Previous work in our unit has shown good correlation between the AVVQ and CEAP, and AVVQ and generic quality of life measures, but poor correlations between VCSS and AVVQ; VCSS and (SQOR-V); and AVVQ and SQOR-V.⁴ Current work being undertaken in our unit has shown excellent correlation between CIVIQ and AVVQ.⁵

Most concerning, however, is the number of patients with C2S disease (symptomatic disease) for whom the authors feel it is appropriate to deny intervention (47% of their cohort). This is in disagreement with extensive work that details the progression of venous disease⁶ and the cost of treating only complicated disease,⁷ not to mention the significant quality of life impairments seen with symptomatic disease⁸ which improve with treatment.⁹ Finally, it has previously been shown that treating uncomplicated venous disease leads to a greater improvement than leaving the disease to progress to permanent damage¹⁰ – the very patients excluded have the most to gain.

Yours Sincerely,

REFERENCES

- 1 Ward A, Abisi S, Braithwaite BD. An online patient completed Aberdeen Varicose Vein Questionnaire can help to guide primary care referrals. *Eur J Vasc Endovasc Surg: Off J Eur Soc Vasc Surg* 2012 Dec 14:1–5.
- 2 Launois R, Reboul-Marty J, Henry B. Construction and validation of a quality of life questionnaire in chronic lower limb venous insufficiency (CIVIQ). *Qual Life Res* 1996 Dec;5(6): 539–54.
- 3 Guex J-J, Zimmet SE, Boussetta S, et al. Construction and validation of a patient-reported outcome dedicated to chronic venous disorders: SQOR-V (specific quality of life and outcome response – venous). *J Mal Vasc* 2007 Jul;32(3):135–47.
- 4 Shepherd AC, Gohel MS, Lim CS, et al. A study to compare disease-specific quality of life with clinical anatomical and hemodynamic assessments in patients with varicose veins. *J Vasc Surg* 2011 Feb;53(2):374–82.
- 5 Kuet ML, Lane TRA, Franklin IJ, et al. A study to compare disease specific quality of life scoring systems in patients with varicose veins. USA: Phoenix; 2013.
- 6 Rabe E, Pannier F, Ko A, et al. Incidence of varicose veins, chronic venous insufficiency, and progression of the disease in the Bonn Vein Study II. *J Vasc Surg* 2010 Mar 1;51(3): 791–1.
- 7 Rabe E, Pannier F. Societal costs of chronic venous disease in CEAP C4, C5, C6 disease. *Phlebology* 2010 Oct 1;25(Suppl. 1):64–7.
- 8 Dunić I, Medenica L, Bobić B, et al. Patients' reported quality of life in chronic venous disease in an outpatient service in Belgrade, Serbia. *Eur J Dermatol* 2009 Jan 1;19(6):616–20.
- 9 Lurie F, Kistner RL. Trends in patient reported outcomes of conservative and surgical treatment of primary chronic venous disease contradict current practices. *Ann Surg* 2011 Aug;254(2):363–7.
- 10 Darvall KAL, Sam RC, Bate GR, et al. Changes in health-related quality of life after ultrasound-guided foam sclerotherapy for great and small saphenous varicose veins. *J Vasc Surg* 2010 Apr;51(4):913–20.

T. Lane*, I. Franklin, A. Davies

Academic Section of Vascular Surgery, Imperial College London, Greater London W6 8RF, United Kingdom

*Corresponding author. Tel.: +44 7832219701; fax: +44 2033117362.

Email-addresses: tristanlane@doctors.net.uk, tristan.lane@nhs.net (T. Lane)

Available online 8 February 2013

© 2013 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.ejvs.2013.01.015>

DOI of original article: <http://dx.doi.org/10.1016/j.ejvs.2012.11.016>

Response to 'Re. An Online Patient Completed Aberdeen Varicose Vein Questionnaire Can Help to Guide Primary Care Referrals'

Messrs Lane, Franklin and Davies have provided additional points for discussion which should be considered if the study is to be repeated. The topic of limiting healthcare is controversial and we are not surprised that our article attracted their response.

We agree that the use of the on-line AVVQ tool will result in patients with C2 disease being denied access to state funded healthcare with the potential outcomes described. It was not our decision to limit access to healthcare but that of the commissioners who set a budget that they could afford. Whether we agree with the concept of limiting the treatment of venous disease to certain groups was not